

普通期水稻定期調査結果(令和7年9月上旬)

25株調査

調査月日：9月1日～3日

調査地点	品種	葉いもち		紋枯病		セジロウンカ			トビイロウンカ				ヒメビロウンカ			コブノメイガ		備考
		発生株率	発病程度	発生株率	発病程度	寄生株率	最高頭数成虫	幼虫の頭数	寄生株率	最高頭数成虫	幼虫の頭数	短翅雌	寄生株率	最高頭数成虫	幼虫の頭数	発生株率	被害程度	
G地区 ①	さがびより	0		0		0			0				0			0		内穎褐変病8%
" ②	夢しずく	0		4	中	16		1	0				4		1	0		
" ③	夢しずく	0		12	中	8		2	8		4	中	8		2	0		
" ④	夢しずく	0		44	中	20	1	2	4		1	中	24		2	0		
H地区 ①	さがびより	0		64	多	16	1	1	0				0			0		
" ②	さがびより	12	少	28	多	20	1	1	8		1	中	0			0		
" ③	ヒノヒカリ	0		8	多	28	1	2	4		1	中	16		2	0		
" ④	さがびより	0		8	中	32	1	2	0				16	1	2	0		
I地区 ①	ヒヨクモチ	4	少	0		4		1					0			44	少	内穎褐変病24%
" ②	ヒヨクモチ	0		0		4		2					0			20	少	
" ③	さがびより	16	少	32	中	4	1						0			0		
" ④	さがびより			12	少	4	1						0			0		
J地区 ①	さがびより	0		0		28	1	1	4		1	老	28	1	1	0		
" ②	さがびより	0		36	少	4	1	1	0				4		1	8	少	
" ③	ヒヨクモチ	0		0		4	1		0				0			0		
" ④	ヒヨクモチ	0		8	少	8	2		0				0			8	少	
K地区 ①	ヒヨクモチ	0		16	少	0			4	1			0			0		
" ②	ヒヨクモチ	0		68	少	0			4	1			0			0		
" ③	ヒヨクモチ	0		12	少	0			0				0			0		
" ④	ヒヨクモチ	0		24	少	4		1	0				0			0		
L地区 ①	ヒヨクモチ	0		0		4		1	0				0			8	少	
" ②	ヒヨクモチ	0		4	少	0			0				0			4	少	
" ③	ヒヨクモチ	0		32	少	4		1	0				0			0		
" ④	ヒヨクモチ	0		0		16	2	2	0				0			0		
M地区 ①	夢しずく	0		12	少	0			0				0			0		もみ枯細菌病4%
" ②	夢しずく	0		0		0			0				0			0		もみ枯細菌病28%
" ③	夢しずく	0		4	少	0			0				0			0		内穎褐変病4%
" ④	夢しずく	0		0		0			0				0			0		もみ枯細菌病4%
N地区 ①	ひなたまる	0		0		0			0				0			0		内穎褐変病8%
" ②	夢しずく	0		4	中	0			0				0			0		
" ③	夢しずく	0		32	中	0			0				0			0		
" ④	夢しずく	0		24	中	0			0				0			0		
O地区 ①	さがびより	0		24	少	12	1	2	0				0			0		
" ②	ヒヨクモチ	0		24	少	4	1		0				0			0		
" ③	ヒヨクモチ	0		8	中	0			0				0			12	少	
" ④	ヒヨクモチ	0		8	中	8	1	3	0				0			8	少	
P地区 ①	ヒヨクモチ	0		0		12		1	0				12		1	48	中	内穎褐変病8%
" ②	さがびより	0		0		16	1	1	0				36		2	0		
" ③	ひなたまる	0		0		12	1	1	0				40	1	3	0		
" ④	ひなたまる	0		0		0			0				0			0		
Q地区 ①	夢しずく	0		0		0			0				0			0		もみ枯細菌病52%
" ②	夢しずく	0		0		12		2	4		1	中	0			0		もみ枯細菌病16%
" ③	さがびより	0		0		0			0				0			0		
" ④	さがびより	0		4	少	0			0				0			0		
R地区 ①	さがびより	0		0		0			0				0			0		内穎褐変病24%
" ②	夢しずく	0		0		0			12		4	中老	12		1	0		もみ枯細菌病4%
" ③	夢しずく	0		0		0			0				0			0		内穎褐変病16%
" ④	夢しずく	0		0		0			28	1	4	中老	0			0		内穎褐変病4%
発生圃場率		6.4		56.3		54.2			22.7				22.9			18.8		
平均発生率		0.7		11.6		6.3			1.8				4.2			3.3		
発生率の平年値		10.4		8.1		4.2			5.1				3.5			15.7		
(平年比)		少		やや多		やや多			並				並			やや少		
発生率の前年値		4.3		11.6		1.3			0.3				2.3			36.5		