

早期水稲定期調査結果(令和8年6月)

25株調査

調査月日：6月16日															
調査地点		葉いもち		紋枯病		セジロウンカ			トビイロウンカ			ヒメビウカ			備考
		発生株率	発病程度	発生株率	発病程度	発生株率	最高頭数 成虫 幼虫	幼虫の齢	発生株率	最高頭数 成虫 幼虫	幼虫の齢	発生株率	最高頭数 成虫 幼虫		
A 地区	①	0		0		0			0			0			
"	②	0		0		4	1	若	0			0			
"	③	0		0		0			0			0			
"	④	0		0		0			0			0			
B 地区	①	0		0		0			0			0			
"	②	0		0		0			0			0			
"	③	0		0		0			0			0			
"	④	0		0		0			0			4		1	
C 地区	①	0		0		16	1	若	0			64	1	5	
"	②	0		0		12	1	若~中	0			0			
"	③	0		0		0			0			0			
"	④	0		0		0			0			0			
発生圃場率		0		0		25.0			0			16.7			
平均発生率		0		0		2.7			0			5.7			
発生率の平年値		1.5		0		6.7			0			8.4			
(平年比)		やや少		並		やや少			並			並			
発生率の前年値		0.7		0		6.0			0			0.7			