

早植え水稻定期調査結果(令和8年8月下旬)

25株調査

調査月日：8月18日

調査地点		葉いもち		紋枯病		セジロウンカ			トビイロウンカ					ヒメトビウンカ			コブノメイガ	フタビコヤガ	備考	
		発生株率	発病程度	発生株率	発病程度	寄生株率	最高頭数 成虫	最高頭数 幼虫	幼虫の齢	寄生株率	最高頭数 成虫	最高頭数 幼虫	幼虫の齢	短翅雌成虫	寄生株率	最高頭数 成虫	最高頭数 幼虫	発生株率		寄生株率
D地区	①	0		0		0				0					0			0	0	
"	②	0		0		0				0					0			0	0	
"	③	0		4	少	0				0					0			4	0	
"	④	4	少	4	少	0				0					0			8	0	
E地区	①	0		0		0				0					0			0	0	ミナミアカメムシ
"	②	0		0		8	1	3	若	0					0			0	0	
"	③	4	少	0		0				0					0			0	0	
"	④	0		0		0				0					0			0	0	
F地区	①	0		0		4	0	1	中	0					0			0	0	イナカメムシ クモヘリカメムシ
"	②	0		0		24	0	1	若中	8	0	1	中		0			0	0	
"	③	0		0		0				0					0			20	0	
"	④	0		0		0				0					0			4	12	
発生圃場率		16.7		16.7		25.0			8.3					0			33.3	8.3		
平均発生率		0.7		0.7		3.0			0.7					0			3.0	1.0		
発生率の平年値		0.9		1.3		6.4			7.3					4.0			3.5	0.8		
(平年比)		並		並		並			並					やや少			並	並		
発生率の前年値		1.7		2.7		1.3			0					0			0	0		